

REPEAT PRESSURES PATIENT INFORMATION SHEET

Measuring your eye pressure

The pressure in one or both of your eyes was higher than normal when it was measured for the first time. This may just have been one abnormally high reading. So your eye pressure needs to be measured again, perhaps twice, to see if the pressure is high each time. Often a different instrument is used for this. When your eye pressure is measured again, the Optometrist will put drops in your eye to numb the surface. Then he or she will use a small instrument to gently touch the front of your eye.

What if all my eye pressure readings are high?

If your eye pressure readings are all above normal, you may have a condition called ocular hypertension (OHT). This means high pressure in the eye. It is important to investigate this further because OHT increases the risk of glaucoma, which can damage your sight. However, most people with OHT never get glaucoma. (There is more about glaucoma at the end of this leaflet.)

What happens next?

Your Optometrist will check for other signs of glaucoma.

If the only sign of glaucoma is that your eye pressure is slightly higher than normal, then taking the measurement again several times will show whether your eye pressure continues to be abnormally high. If it is, then your optometrist will arrange for you to visit a Hospital eye clinic. However, if the repeat test shows your eye pressures are within the normal range, then you simply need to have further regular eye examinations with your optometrist.

Will I need more tests?

If the pressure readings are high after repeating the measurement, your Optometrist will make an appointment for you to be seen on another day so that these tests may be repeated. Repeating these measurements will allow your optometrist to make a clearer assessment of your eye pressures.

If the repeated eye pressure test gives measurements that are similar to the original findings, this suggests that there is a strong suspicion that you may have ocular hypertension. Your Optometrist will then arrange for you to be seen at a hospital eye clinic.

If the repeated tests are not similar, but show findings that are within normal ranges, you will need only to have regular examinations with your Optometrist. They will advise you if you need to be seen more frequently.

What happens if I'm referred to the Hospital eye clinic?

If you are referred to the Hospital, your Optometrist will send a referral directly to the Hospital and also inform your GP.

You should receive an offer of an appointment from the Hospital within 6 weeks of referral. If you do not receive a Hospital appointment letter within 6 weeks, then please call the Central Booking Office to enquire about your referral on any of the following telephone numbers:

0300 422 Followed by 5973 or 5975 or 6871 or 6878 or 6950 or 5997

Failing to attend appointments to repeat measurements

Your Optometrist will arrange an appointment for you to have the pressure test repeated if the results indicate it is necessary. If you fail to attend this appointment, and the repeat readings therefore are not achieved, your Optometrist will refer you to your GP.

What is Glaucoma?

Glaucoma is a group of eye conditions in which the pressure of the fluid inside the eye damages optic nerve (connecting the eye to the brain).

The common form of glaucoma is the chronic form. This is usually very slow to progress. If it is not treated the damage reduces the peripheral (side) vision and can eventually result in tunnel vision – like looking down a long tube. Most people are not aware of any changes in the early stages of chronic glaucoma. The best way to catch it early is to have regular eye examinations. Chronic glaucoma can be treated effectively, usually with drops.

There is a much rarer, acute form of glaucoma, where the pressure in the eye rises rapidly.

This needs immediate treatment in hospital to avoid permanent damage to your sight. This acute form of glaucoma is often very painful; the eye may become red, you may feel sick and vomit. If this happens you should go to the Accident and Emergency department immediately. In the early stages there may be rainbow coloured rings around lights or it may be like looking through a mist; the eyes may ache and these symptoms can sometimes come and go. It is important to act quickly; seek advice as soon as possible to prevent sight loss.

If you need any more advice or explanation, please ask you Optometrist.

About your records

Primary Eyecare Gloucestershire collects the details of these repeated measurements and keeps a record of them secure on a web based system. If you are referred to other health professionals, these details may be shared with them to ensure you receive the best care. Some information is also used to check how the service to the patient is working, make improvements and administer finances.

Personal details are not shared with any other groups or individuals.

If you do not wish Primary Eyecare Gloucestershire to hold and use your details in this way, please tell the Optometrist who will arrange for the details to be removed from the computer system.

Patient Satisfaction

We would like to hear of your experience with this service. Therefore, we would be most grateful if you would complete the short survey that the Optometrist will give you. Your feedback is valuable and will help us continually monitor and develop the service.